

File 347:JAPIO OCT 1976-1996/DEC.(UPDATED 970505)

(c) 1997 JPO & JAPIO

*File 347: Records current through Kokai Number 08-340700

Set Items Description

?t 05208491/9

05208491/9

DIALOG(R)File 347:JAPIO

(c) 1997 JPO & JAPIO. All rts. reserv.

05208491

PRODUCTION OF TAXANE-TYPE DITERPENE

PUB. NO.: 08-163991 [JP 8163991 A]

PUBLISHED: June 25, 1996 (19960625)

INVENTOR(s): YUKIMUNE TAKAHITO

HARA YASUHIRO

APPLICANT(s): MITSUI PETROCHEM IND LTD [000588] (A Japanese Company or Corporation), JP (Japan)

APPL. NO.: 06-312258 [JP 94312258]

FILED: December 15, 1994 (19941215)

INTL CLASS: [6] C12P-013/02; A61K-031/21; A61K-031/335; C12P-013/02; C12R-001/91

JAPIO CLASS: 14.5 (ORGANIC CHEMISTRY -- Microorganism Industry); 14.4 (ORGANIC CHEMISTRY -- Medicine)

JAPIO KEYWORD:R051 (PHARMACEUTICALS -- Anti-cancer Agents)

ABSTRACT

PURPOSE: To obtain a taxane-type diterpene compound useful for a medicine, etc., of ovarian cancer, breast cancer and lung cancer, etc., in a high productivity by culturing tissue or cell, etc., of a plant producing a taxane-type diterpene in a medium containing fatty acids and recovering a product from the resultant cultured material.

CONSTITUTION: A tissue and/or cell of a plant (e.g. *Taxus brevifolia* NUTT) producing taxane-type diterpene (e.g. taxol, et.) is cultured in a medium containing 10-22C fatty acids such as oleic acid, linoleic acid, linolenic acid, arachidonic acid, capric acid, decenoic acid, lauric acid, dodecenoic acid, myristic acid, palmitic acid, stearic acid, eicosapentaenoic acid or docosaheptaenoic acid in a concentration of 0.01-1000.mu.M. The object taxane-type diterpene useful for a remedy of ovarian cancer, breast cancer and lung cancer, etc., is obtained in a high productivity by recovering the taxane-type diterpene from the resultant cultured material.

?t 05071510/9

05071510/9

DIALOG(R)File 347:JAPIO

(c) 1997 JPO & JAPIO. All rts. reserv.

BT
B10